

Matched Asymptotic Expansions of the Eigenvalues of a 3-D boundary-value problem relative to two cavities linked by a hole of small size

M'BAREK FARES¹, ABDELKADER TIZAOU², SÉBASTIEN TORDEUX²

¹Electromagnetism and Acoustics, CERFACS, 42 Avenue Gaspard Coriolis, F-31100 Toulouse

²Toulouse University, INSA-Toulouse, Mathematical Institute of Toulouse (UMR-CNRS 5219), 135 avenue de Rangueil, F-31077 Toulouse, France, stordeux@insa-toulouse.fr

In a lot of physical problems, the boundary of the computational domain is perforated. This configuration can lead to numerical difficulties when the diameter of the holes are really smaller than the other characteristic lengths. Indeed, it can be very costly to compute a sharp numerical approximation of the solution of such problems for two main reasons: With a standard method like finite elements or finite differences, a refined mesh cannot be avoided in the neighborhood of the hole; the mesh generation of a perforated structure can be a hard task.

Many authors have studied the effect of the perforation of the boundaries both from the theoretical and the numerical point of views, see for example [3–5]. However, less results have been obtained for the eigenvalue problem in the case of a three dimensional domain.

In [2], Gadyl'shin considered a two dimensional domain consisting of two domains linked by a small hole. He derived a complete asymptotic expansion of the scattering frequencies of the Laplacian operator equipped with Dirichlet boundary condition. In [1], these results were extended to the eigenvalues and eigenvectors of an elliptic operator with varying coefficients. In this talk, we are interested to a three dimensional configuration with varying coefficients and Neumann boundary condition.

References

- [1] A. BENDALI, A. HUARD, A. TIZAOU, S. TORDEUX AND J. P. VILA, *Asymptotic Expansions of the Eigenvalues of a 2-D boundary-value problem relative to two cavities linked by a hole of small size*, C. R. Acad. Sci. Paris, Mathmatiques, 347 (2009), pp. 1147-1152.
- [2] R. R. GADYL'SHIN, *Surface potentials and the method of matching asymptotic expansions in the Helmholtz resonator problem*, (Russian) Algebra i Analiz 4 (1992), no. 2, 88–115, translation in St. Petersburg Math. J. 4 (1993), no. 2, 273–296.
- [3] J. SANCHEZ-HUBERT AND E. SÁNCHEZ-PALENCIA, *Acoustic fluid flow through holes and permeability of perforated walls*, J. Math. Anal. Appl., 87 (1982), pp. 427–453.
- [4] A. TAFLOV, K. UMASHANKER, B. BECKER, F. HARFOUSH, AND K. S. YEE, *Detailed fdtd analysis of electromagnetic fields penetrating narrow slots and lapped joints in thick conducting screens*, IEEE Trans Antenna and Propagation, 36 (1988), pp. 2472-57.
- [5] E. O. TUCK, *Matching problems involving flow through small holes*, in Advances in applied mechanics, Vol. 15, Academic Press, New York, (1975), pp. 89–158.